OpenEvidence has signed content agreements with **NEJM**, **JAMA**, **NCCN**, **Wiley**, **Cochrane**, and more.

OpenEvidence

This conversation has been affirmatively and intentionally made public by the user who created it.

Summarize and support complications r/t prehospital needle chest decompression.

Analyzed query, searched for evidence

Prehospital needle chest decompression (PHND) carries a **complication rate of approximately 2–14%** and a **procedural failure rate of 39–76%**, making it a procedure with significant limitations despite its life-saving potential. JAMA Surg + 3

Specific Complications

- **Iatrogenic pneumothorax or hemothorax** — creation of a new pneumothorax or hemothorax in patients who did not have one, or worsening of existing injury. At least **39% of patients** in one large series did not have a tension pneumothorax at all, meaning the procedure was performed unnecessarily. American Surg...
- **Cardiac injury** — a rare but devastating complication. In one series, a needle was inserted into the heart, requiring cardiac surgery and resulting in anoxic brain injury. Anatomic studies show that using an 8-cm catheter at the 5th intercostal space anterior axillary line places the heart at risk of iatrogenic injury in **42% of patients**. Injury + 1
- **Subcutaneous hematoma** — localized bleeding at the insertion site. JAMA Surg
- **Catheter dislodgement or kinking** — leading to loss of decompression and potential hemodynamic compromise. JAMA Surg
- **Iatrogenic injury (broadly)** — one study found iatrogenic injury rates of **16.7–28.2%** depending on the era studied, with only modest improvement after protocol and curriculum changes. American Jour...

Procedural Failure

The most significant "complication" is **failure to reach the pleural space**. A 10-year retrospective analysis found that **39–76% of needle decompression attempts failed** to reach the pleural cavity, depending on the radiologic method used to assess success.

American Surg... Failure correlates strongly with increased chest wall thickness, and injured chest walls are significantly thicker than uninjured ones at both the 2nd ICS midclavicular and 5th ICS anterior axillary sites. American Surg... Only **30% of patients** in another large series demonstrated a positive clinical response (improved SpO2, SBP, breath sounds, or ROSC), though this improved to **48%** when patients already in cardiac arrest were excluded. Injury

Misapplication and Patient Selection

A major contributor to poor outcomes is **inappropriate patient selection**. The majority of procedures are performed based on subjective indicators (absent/reduced breath sounds, ongoing CPR) rather than objective criteria such as confirmed hypoxia. Injury The NAEMSP position statement highlights frequent misapplication and low rates of successful decompression as key concerns. Prehospital Em... One study found that proper patient selection — defined as the presence of decompensated shock, respiratory distress, and diminished breath sounds — was present in only **23–28% of cases**. American Jour...

Mortality Context

Despite these complications, PHND was associated with a **25% decrease in odds of 24-hour mortality** compared with emergent tube thoracostomy at the trauma center in a large JAMA Surgery cohort study, suggesting the procedure remains beneficial when appropriately applied. JN JAMA Surg Reported mortality after PHND ranges from **12–79%**, with rates universally lower in patients not presenting in traumatic arrest. JN JAMA Surg

Emerging Solutions

Recent evidence suggests that **colorimetric capnography** as an adjunct during needle decompression may improve initial success rates (82% vs. 66%) and reduce the need for repeat procedures or conversion to finger thoracostomy. American Jour... Prehospital point-of-care ultrasound may also improve patient selection. JN JAMA Surg The NAEMSP and others advocate for incorporating objective criteria and standardized protocols to reduce variability and improve outcomes. Injury + 1

Would you like to explore the comparative evidence between needle thoracostomy and finger thoracostomy as alternative prehospital decompression techniques?

References